

[Skip to main content](#)

Zarontin (Ethosuximide)

Zuletzt aktualisiert: 30. November 2017

Allergic reaction. Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). **Body As A Whole:**

Gastrointestinal symptoms occur frequently and include anorexia, vague gastric upset, nausea and vomiting, cramps, epigastric and **Gastrointestinal System:** abdominal pain, weight loss, and diarrhea. There have been reports of gum hypertrophy and swelling of the tongue.

Hemopoietic complications associated with the administration of ethosuximide have included leukopenia, agranulocytosis, pancytopenia, with **Hemopoietic System:** or without bone marrow suppression, and eosinophilia.

Neurologic and sensory reactions reported during therapy with ethosuximide have included drowsiness, headache, dizziness, euphoria, hiccups, **Nervous System:** irritability, hyperactivity, lethargy, fatigue, and ataxia. Psychiatric or psychological aberrations associated with ethosuximide administration have included disturbances of sleep, night terrors, inability to concentrate, and aggressiveness. These effects may be noted particularly in patients who have previously exhibited psychological abnormalities. There have been rare reports of paranoid psychosis, increased libido, and increased state of depression with overt suicidal intentions.

Dermatologic manifestations which have occurred with the administration of ethosuximide have included urticaria, pruritic erythematous **Integumentary System:** rashes, and hirsutism.

Myopia. **Special Senses:**

Vaginal bleeding, microscopic hematuria. **Genitourinary System:**