

[Skip to main content](#)

Colchicine Capsules (Mitigare)

??? ??????: 30 ?????2/?????? 2017

Gastrointestinal disorders are the most common adverse reactions with colchicine. They are often the first signs of toxicity and may indicate that the colchicine dose needs to be reduced or therapy stopped. These include diarrhea, nausea, vomiting, and abdominal pain.

] **WARNINGS AND PRECAUTIONS** Colchicine has been reported to cause neuromuscular toxicity, which may present as muscle pain or weakness. [see

Toxic manifestations associated with colchicine include myelosuppression, disseminated intravascular coagulation, and injury to cells in the renal, hepatic,] **OVERDOSAGE** circulatory, and central nervous system. These most often occur with excessive accumulation or overdose [see

The following reactions have been reported with colchicine. These have been generally reversible by interrupting treatment or lowering the dose of colchicine:

abdominal cramping, abdominal pain, diarrhea, lactose intolerance, nausea, vomiting **Digestive:**

sensory motor neuropathy **Neurological:**

alopecia, maculopapular rash, purpura, rash **Dermatological:**

leukopenia, granulocytopenia, thrombocytopenia, pancytopenia, aplastic anemia **Hematological:**

elevated AST, elevated ALT **Hepatobiliary:**

myopathy, elevated CPK, myotonia, muscle weakness, muscle pain, rhabdomyolysis **Musculoskeletal:**

azoospermia, oligospermia **Reproductive:**