

[Skip to main content](#)

Cyclessa (Desogestrel Ethinyl Estradiol Tablets)

??? ??????: 30 ?????2/????? 2017

section): **WARNINGS** An increased risk of the following serious adverse reactions has been associated with the use of oral contraceptives (see

- Cerebral hemorrhage
- Cerebral thrombosis
- Hypertension
- Gallbladder disease
- Hepatic adenomas or benign liver tumors
- Thrombophlebitis and venous thrombosis with or without embolism
- Arterial thromboembolism
- Pulmonary embolism
- Myocardial infarction

There is evidence of an association between the following conditions and the use of oral contraceptives:

- Retinal thrombosis
- Mesenteric thrombosis

The following adverse reactions have been reported in patients receiving oral contraceptives and are believed to be drug-related:

- Change in weight or appetite (increase or decrease)
- Change in cervical ectropion and secretion
- Possible diminution in lactation when given immediately postpartum
- Cholestatic jaundice
- Migraine headache
- Rash (allergic)
- Mood changes, including depression
- Vaginitis, including candidiasis
- Change in corneal curvature (steepening)
- Intolerance to contact lenses
- Exacerbation of systemic lupus erythematosus
- Exacerbation of chorea
- Anaphylactic/anaphylactoid reactions, including urticaria, angioedema, and severe reactions with respiratory and circulatory symptoms
- Nausea
- Vomiting
- Gastrointestinal symptoms (such as abdominal pain, cramps and bloating)
- Breakthrough bleeding
- Spotting
- Change in menstrual flow
- Amenorrhea
- Temporary infertility after discontinuation of treatment
- Edema/fluid retention
- Melasma/chloasma which may persist
- Breast changes: tenderness, pain, enlargement, and secretion
- Decrease in serum folate levels
- Exacerbation of porphyria
- Aggravation of varicose veins

The following adverse reactions have been reported in users of oral contraceptives and the association has been neither confirmed nor refuted:

- Erythema nodosum
- Hemorrhagic eruption
- Impaired renal function
- Hemolytic uremic syndrome
 - Acne
 - Changes in libido
 - Colitis
 - Budd-Chiari Syndrome
- Optic neuritis, which may lead to partial or complete loss of vision
- Pre-menstrual syndrome
 - Cataracts
- Cystitis-like syndrome
 - Headache
 - Nervousness
 - Dizziness
 - Hirsutism
 - Loss of scalp hair
- Erythema multiforme
 - Dysmenorrhea
 - Pancreatitis