

[Skip to main content](#)

## Diuril (Chlorothiazide)

??? ??????: 30 ?????2/????? 2017

The following adverse reactions have been reported and, within each category, are listed in order of decreasing severity.

Weakness. **Body as a Whole:**

Hypotension including orthostatic hypotension (may be aggravated by alcohol, barbiturates, narcotics or antihypertensive drugs). **Cardiovascular:**

Pancreatitis, jaundice (intrahepatic cholestatic jaundice), diarrhea, vomiting, sialadenitis, cramping, constipation, gastric irritation, nausea, anorexia. **Digestive:**

Aplastic anemia, agranulocytosis, leukopenia, hemolytic anemia, thrombocytopenia. Hypersensitivity: Anaphylactic reactions, necrotizing angitis **Hematologic:**  
(vasculitis and cutaneous vasculitis), respiratory distress including pneumonitis and pulmonary edema, photosensitivity, fever, urticaria, rash, purpura.

), hyperglycemia, glycosuria, hyperuricemia. **PRECAUTIONS** Electrolyte imbalance (see **Metabolic:**

Muscle spasm. **Musculoskeletal:**

Vertigo, paresthesias, dizziness, headache, restlessness. **Nervous System/Psychiatric:**

.) **WARNINGS** Renal failure, renal dysfunction, interstitial nephritis. (See **Renal:**

Erythema multiforme including Stevens-Johnson syndrome, exfoliative dermatitis including toxic epidermal necrolysis, alopecia. **Skin:**

Transient blurred vision, xanthopsia. **Special Senses:**

Impotence. **Urogenital:**

Whenever adverse reactions are moderate or severe, thiazide dosage should be reduced or therapy withdrawn.