

[Skip to main content](#)

Dyazide (Hydrochlorothiazide and Triamterene)

??? ??????: 30 ?????2/????? 2017

Adverse effects are listed in decreasing order of severity.

: Anaphylaxis, rash, urticaria, subacute cutaneous lupus erythematosus-like reactions, photosensitivity. **Hypersensitivity**

: Arrhythmia, postural hypotension. **Cardiovascular**

: Diabetes mellitus, hyperkalemia, hypokalemia, hyponatremia, acidosis, hypercalcemia, hyperglycemia, glycosuria, hyperuricemia, hypochloremia. **Metabolic**

: Jaundice and/or liver enzyme abnormalities, pancreatitis, nausea and vomiting, diarrhea, constipation, abdominal pain. **Gastrointestinal**

: Acute renal failure (one case of irreversible renal failure has been reported), interstitial nephritis, renal stones composed primarily of triamterene, elevated **Renal** BUN, and serum creatinine, abnormal urinary sediment.

: Leukopenia, thrombocytopenia and purpura, megaloblastic anemia. **Hematologic**

: Muscle cramps. **Musculoskeletal**

: Weakness, fatigue, dizziness, headache, dry mouth. **Central Nervous System**

: Impotence, sialadenitis. **Miscellaneous**

Thiazides alone have been shown to cause the following additional adverse reactions:

: Paresthesias, vertigo. **Central Nervous System**

: Xanthopsia, transient blurred vision. **Ophthalmic**

: Allergic pneumonitis, pulmonary edema, respiratory distress. **Respiratory**

: Necrotizing vasculitis, exacerbation of lupus. **Other**

: Aplastic anemia, agranulocytosis, hemolytic anemia. **Hematologic**

: Thrombocytopenia and pancreatitis-rarely, in newborns whose mothers have received thiazides during pregnancy. **Neonate and infancy**

: Erythema multiforme including Stevens-Johnson syndrome, exfoliative dermatitis including toxic epidermal necrolysis. **Skin**