

[Skip to main content](#)

EEMT (Esterified Estrogens and Methyltestosterone Tablets)

??? ??????: 30 ?????2/?????? 2017

Associated with Estrogens

WARNINGS regarding induction of neoplasia, adverse effects on the fetus, increased incidence of gallbladder disease, and adverse effects similar (See). The following additional adverse reactions have been reported with estrogenic therapy, including **to those of oral contraceptives, including thromboembolism** oral contraceptives:

Genitourinary system

Breakthrough bleeding, spotting, change in menstrual flow.
Dysmenorrhea.
Premenstrual-like syndrome.
Amenorrhea during and after treatment.
Increase in size of uterine fibromyomata.
Vaginal candidiasis.
Change in cervical erosion and in degree of cervical secretion.
Cystitis-like syndrome.

Breasts

Tenderness, enlargement, secretion.

Gastrointestinal

Nausea, vomiting.
Abdominal cramps, bloating.
Cholestatic jaundice.

Skin

Chloasma or melasma which may persist when drug is discontinued.
Erythema multiforme.
Erythema nodosum.
Hemorrhagic eruption.
Loss of scalp hair.
Hirsutism.

Eyes

Steepening of corneal curvature.
Intolerance to contact lenses.

CNS

Headache, migraine, dizziness.
Mental depression.
Chorea.

Miscellaneous

Increase or decrease in weight.
Reduced carbohydrate tolerance.

Aggravation of porphyria.
Edema.
Changes in libido.

Associated with Methyltestosterone

Endocrine And Urogenital

The most common side effects of androgen therapy are amenorrhea and other menstrual irregularities, inhibition of gonadotropin secretion, **Female:** .1
and virilization, including deepening of the voice and clitoral enlargement. The latter usually is not reversible after androgens are discontinued. When
administered to a pregnant woman androgens cause virilization of external genitalia of the female fetus.
Hirsutism, male pattern of baldness, and acne. **Skin and Appendages:** .2
Retention of sodium, chloride, water, potassium, calcium, and inorganic phosphates. **Fluid and Electrolyte Disturbances:** .3
Nausea, cholestatic jaundice, alterations in liver function test, rarely hepatocellular neoplasms, and peliosis hepatis (see **Gastrointestinal:** .4
)**.WARNINGS**
Suppression of clotting factors, II, V, VII, and X, bleeding in patients on concomitant anticoagulant therapy, and polycythemia. **Hematologic:** .5
Increased or decreased libido, headache, anxiety, depression, and generalized paresthesia. **Nervous System:** .6
Increased serum cholesterol. **Metabolic:** .7
Inflammation and pain at site of intramuscular injection or subcutaneous implantation of testosterone containing pellets, stomatitis with **Miscellaneous:** .8
buccal preparations, and rarely anaphylactoid reactions.