

Anthocyanin pigments: Structure and biological importance

Rana Nassour^{1*}, Abdulkarim Ayash¹, Kanaan Al-Tameemi²

¹Department of Basic Sciences, Faculty of Pharmacy, Al-Andalus University for Medical Sciences, Tartous, Syria.

²Department of Microbiology, Faculty of Pharmacy, Al-Andalus University for Medical Sciences, Tartous, Syria.

*Corresponding author: ranahn1985@gmail.com

ABSTRACT

Anthocyanins are coloured water-soluble pigments representing one of the major subclasses of compounds. They rarely exist in nature as free aglycons, instead, they attach to one or more sugar moieties. Anthocyanins are found within different plant organs; flowers, leaves, fruits, roots, tubers and grains. They appear in different attractive colours depending on their structure, pH, and other factors.

These compounds gained a lot of attention in the last few years as food colourants replacing chemical dyes, besides their role in enhancing plant tolerance against many abiotic stresses such as salinity, drought, excessive light, ultraviolet radiation and cold stress. Besides, previous studies demonstrated the importance of anthocyanins in human health and their protective properties against chronic diseases. Hence, this review focuses on anthocyanins as one of the most important pigments having beneficial roles in health for plants and humans.

KEY WORDS: Anthocyanins, colourants, plant stress, antioxidants, antibacterial, anti-inflammatory, anticancer, anti-diabetic, anti-obesity, neurodegenerative diseases, cardiovascular diseases, ophthalmology.

1. INTRODUCTION

Anthocyanins (Greek anthos: flower and kyaneos: dark blue) represent a subclass of the phenolic compounds (Delgado-Vargas, 2000). They are water-soluble glycosides of anthocyanidins, which are largely responsible for the attractive pale yellow, orange, red, magenta, violet and blue colour of a wide range of plant tissues, principally flowers, leaves and fruits, besides storage organs, roots, tubers, stems and grains (Chemler, 2009; Martin, 2017). They are ubiquitous in higher plants (occurring in more than 30 families), but normally are absent in liverworts, algae, and other lower plants, even though some anthocyanins have been identified in mosses and ferns (Delgado-Vargas, 2000).

In general, anthocyanins are non-photosynthetic pigments synthesized in the cytoplasm and stored in the vacuolar lumen of epidermal cells (Andersen and Jordheim, 2006; Chemler, 2009; Chanoca, 2015; Passeri, 2016).

Within the main vacuole, different forms of anthocyanins accumulation were observed and called by several names reflecting their diversity. Most cells exhibit an evenly coloured anthocyanin vacuolar sap, but there are now many reported cases of regular or irregular shaped pigmented bodies that have been called anthocyanin vacuolar inclusions (AVIs) (Grotewold and Davies, 2008).

It's worth mentioning that cytoplasmic anthocyanin enters the vacuole through microautophagy, which includes three stages, (Figure.1) (Chanoca, 2015; Oku and Sakai, 2018):

- Stage I: The anthocyanins aggregates (AA) associate with the outer surface of the tonoplast and become surrounded by a double membrane. The tonoplast binds closely to the surface of the AA and lines its internal holes. After that, the distant domains of the tonoplast edges fuse.
- Stage II: The AA's two membranes start to separate, and the bulges in between are filled with the vacuolar lumen.
- Stage III: The two membranes separate, releasing the new AVI into the vacuolar lumen. AVI is now surrounded by one membrane derived from the tonoplast.




Figure.1. AVI formation by micro autophagy (Chanoca, 2015)

Based on the extensive studies on AVIs, AVIs have two distinct types. One of them has spherical shaped with a liquid behavioural, mobile and usually coalesce into bigger bodies. While the second type of AVIs has a more 'fibrous' rigid and insoluble structure, and they have been mostly studied in petal cells of many plants (Grotewold and Davies, 2008; Deroles, 2009).

The anthocyanin molecule consists of an anthocyanidin "core" with an attached sugar moiety (Figure.2). They vary in the number and position of hydroxyl and methoxyl groups attached to anthocyanidin. Therefore, although there are nearly 25 naturally occurring anthocyanidins, more than 700 anthocyanins derivatives have been identified to date (Mortensen, 2006; Chu, 2011; Wallace and Giusti, 2015; 2019).




Figure.2. Basic structure of anthocyanidins (Zhao, 2018)

The most prevalent anthocyanidins in nature are pelargonidin (Pg), cyanidin (Cy), peonidin (Pn), delphinidin (Dp), petunidin (Pt) and malvidin (Mv) (Vermerris and Nicholson, 2006; Latti, 2009; Chu, 2011; Khoo, 2017; Wallace and Giusti, 2019):

- Pelargonidin (Pg) appears in a unique orange salmon colour. It can be found in various plants like strawberry, blueberry, banana, red radish and potato (Deroles, 2009; Bueno, 2012; Fang, 2015).
- Cyanidin (Cy) usually appears in magenta and crimson colours. The major sources of Cy are apples, blackberry, black raspberries, elderberry, mulberry, bilberry, gooseberry, peach, pear, fig, cherry, red onion, red cabbage, blood orange, plum, grape, red sweet potatoes, potatoes, strawberry and purple carrot (Deroles, 2009; Usenik, 2009; Chu, 2011; Bueno, 2012; Fang, 2015).
- Peonidin (Pn) has a magenta colour. It can be found in many fruits like mango, grape, potatoes, sweet potatoes and plum (Deroles, 2009; Usenik, 2009; Bueno, 2012; Fang, 2015).
- Delphinidin (Dp) normally is found in different colours; purple, mauve and blue. The blue hue of flowers is due to this pigment (Khoo, 2017). The main sources for it are passion fruit, eggplant, green bean, pomegranate, blueberry, bilberry and grape (Deroles, 2009; Chu, 2011; Bueno, 2012; Fang, 2015).
- Petunidin (Pt) has a purple colour. Pt is found in blueberry, bilberry and grape (Deroles, 2009; Chu, 2011; Bueno, 2012; Fang, 2015).
- Malvidin (Mv) has a purple colour. It is primarily responsible for the colour of bilberry, blueberry and red grape (Deroles, 2009; Chu, 2011; Bueno, 2012; Fang, 2015).

Anthocyanidins rarely occur in nature as free aglycones due to their instability, so when they are glycosylated with one or more sugar molecules they are called anthocyanins. The most common sugar that binds to anthocyanidins is glucose, but there are other sugars involved in anthocyanins formation, such as arabinose, galactose, xylose, and rhamnose (Vermerris and Nicholson, 2006; Tomic, 2017).

During glycosylation, if one sugar bonds with anthocyanidin, it will mainly attach at the 3-position on the C-ring to form the 3-glycoside, and if a second sugar is present, it will almost always attach at the 5-position on the A-ring and form di-glycoside. Yet, there are a few rare exceptions with 3,7-substitutions or trisaccharide forms of anthocyanins (Harborne and Williams, 2001; Wu and Prior, 2005; Prior and Wu, 2006; Vermerris and Nicholson, 2006; Latti, 2009).

The sugar moiety may be acylated by aromatic acids, mainly hydroxycinnamic acids (such as caffeic, ferulic, p-coumaric and sinapic acids) and sometimes by aliphatic acids (particularly malonic and acetic acids). These acyl moieties are usually linked to the sugar at the 3-position on the C-ring (Pereira, 2009; Bueno, 2012).

The Colour of Anthocyanins: Intensity and type of anthocyanin colour are affected by the number of hydroxyl and methoxyl groups in their structure. If it contains more hydroxyl groups (OH), the colour goes toward a more bluish shade. On the other hand, the redness increases if it includes more methoxyl groups (OCH₃) (Mortensen, 2006; Tanaka, 2008; Bueno, 2012; Martin, 2017).

Similarly, the colour of anthocyanin depends on pH, due to their ionic nature. In acidic conditions, some of the anthocyanins appear red. They have a purple hue in neutral pH, while the colour changes to blue in an increasing pH condition (Khoo, 2017).

However, the colour of anthocyanin depends on other factors, including metal ions present and the extent of anthocyanin glycosylation and acylation, beside the combinations of several anthocyanidins and other pigmentation (Gould, 2002; Andersen and Jordheim, 2006; Vermerris and Nicholson, 2006). Finally, storage may also affect anthocyanin stability leading to colour change (Santos-Buelga, 2014).

Anthocyanins as Antioxidants: Living cells produce byproducts during metabolism in the form of reactive oxygen species (ROS) and free radicals, under normal and stressed conditions. ROS are a group of reactive molecules derived from molecular oxygen, such as superoxide (O_2^-), singlet oxygen (1O_2), hydrogen peroxide (H_2O_2) and hydroxyl radical (OH \cdot). They can induce cellular damage when excessively produced (Martin, 2017).

Anthocyanins and anthocyanidins have a higher antioxidant property compared to other flavonoids, due to their special chemical structure. The antioxidant capacity of these compounds can be attributed to chelate metal ions involved in free radicals production, thereby reducing metal-induced peroxidation (Dai, 2012; Martin, 2017). Additionally, their positive charge, number and position of hydroxyl and methoxyl groups, the presence of electron-donating and electron-withdrawing substituents made anthocyanins very effective donors of hydrogen to ROS and free radicals, thereby detoxifying them and preventing further radical formation. This effect protects the important biomolecules (proteins, lipids and DNA) from oxidative damage, which leads to ageing and various diseases (Pojer, 2013; Martin et al., 2017).

The antioxidant activity of anthocyanins increases with the number of hydroxyl groups in the B-ring (Liu, 2018). On the other hand, glycosylation of anthocyanins decreases scavenger activity as compared with their aglycones, because it minimizes the hydrogen donating and metal chelating abilities (Wang and Stoner 2008; Pojer, 2013; Liu, 2018). Moreover, different attached sugars influence antioxidant activity differently (Sadilova, 2006; Liu, 2018).

Anthocyanins can scavenge free radicals through two hypothesized pathways. The first pathway is the attack of OH group(s) of the B-ring, and the second is the attack of oxonium ion on the C-ring. Some of them are considered among the strongest antioxidants via adopting both pathways (Gauljac, 1999; Pojer, 2013; Khoo, 2017).

Most of the widely distributed anthocyanidins and anthocyanins show more scavenging activity than that of the well-known strong antioxidants, for instance, the cyanidin has an antioxidant capacity up to 4.4 times greater than those of ascorbic acid and the vitamin E analogue (Gould, 2002).

The Importance of Anthocyanins: Anthocyanins are believed to play vital roles in plants, animals and human lives. As a major subclass of secondary metabolites in plants commonly consumed as food, they are important in the food industry and human nutrition. Anthocyanins in the flower petals attract insects and birds, so they help in cross-pollination. Similarly, anthocyanins in the colourful skins of fruits attract herbivorous animals that may eat them and disperse the seeds (Hirsch and Martins, 2015). Anthocyanins are important as natural colourants. Furthermore, they possess many functions in plants exposed to abiotic stresses as well as in human health promotion.

Anthocyanins as natural colourants: Anthocyanins are used as food colourants instead of synthetic colourants, as they are safe to be consumed even at relatively high doses (Chemler, 2009; Khoo, 2017).

Different hues can be achieved, depending on anthocyanin chemical structure of anthocyanin, the extraction method and pH of the food matrix. Anthocyanin acylation usually improves colour and pigment stability (Giusti and Wrolstad, 2003; Wallace and Giusti, 2019). For example, acylated pelargonidin derivatives from radish and potato can give an attractive red colour under acidic conditions and acylated cyanidin derivatives from black carrots or cabbage provide hues ranging from deep red to purple, depending on pH (Giusti and Wrolstad, 2003).

Anthocyanin-derived colourants are classified as one of the nine accepted natural colourants under classification E 163 in the European Union (Wrolstad, 2004; Bueno, 2012). However, in 2013, a group of scientific experts for European Food Safety Authority inferred that anthocyanins from various natural sources have been inadequately characterized by safety and toxicological studies to approve their use as food additives (European Food Safety Authority, 2013).

While in the United States, they can be used as natural colourants without certifications (Code of Federal Regulations, 2020). They are also considered appropriate for food use in Asian, Central American, and South American countries; with some differences in their sources and permitted a degree of purity (Wrolstad and Culver, 2012).

Anthocyanins and plant stresses: There is a piece of increasing evidence that anthocyanins have a role to play in the physiological survival of plants under different abiotic stresses, particularly when they are located at the upper surface of the leaf or in the epidermal cells (Gould, 2002; Andersen and Jordheim, 2006; Manetas, 2006; Passeri, 2016).

Cold stress: Low temperatures reduce membrane fluidity, enzymes activities (including the most important one in photosynthesis; RubisCO), and stomatal conductance. Consequently, the photosynthetic rate declines, which induce ROS generation. As toxic substances, ROS attack cellular biomolecules, destroy bio-membranes and accelerate cell damage (Schulz, 2016; Zhang, 2019). Low temperature stimulates anthocyanin biosynthesis by up-regulating the

expression of anthocyanin biosynthetic genes which in turn increase the anthocyanins accumulation (Ahmed, 2015; Schulz, 2015; Schulz, 2016; Zhang, 2019; He, 2020). The main purpose of this accumulation is to increase antioxidant capacity and reduce oxidative stress (Ubi, 2006; Ahmed, 2015; Zhang, 2019).

Salt stress: Anthocyanins content increases under salt stress in many plants (Eryilmaz, 2006; Chakovari, 2015; Kovinich, 2015; Chunthaburee, 2016; Kielkowska, 2019). This increment helps plants to maintain turgidity and water uptake, which is essential for osmo protectant (Close and Beadle, 2003; Iseri, 2015). Besides, the presence of phenyl groups in these compounds induces salt-tolerance by binding with toxic ions, thereby protecting the cells from oxidative damage (Chunthaburee, 2016).

Drought stress: Anthocyanins can mitigate water stress (drought) in different plants, according to previous studies (Sperdouli and Moustakas, 2012; Nakabayashi, 2014; Shoeva, 2017; Kebbas, 2018; Omid, 2018). These compounds could play a dual role under drought stress as osmoregulatory and antioxidants, which allow anthocyanins leaves to tolerate suboptimal water levels (Chalker-Scott, 2002; Kebbas, 2018; Omid, 2018).

High light stress: Foliar anthocyanins concentrations increase in many plants under excessive light condition, acting as photo-protectants (Steyn, 2002; Zhang, 2010; Kovinich, 2015; Trojak and Skowron, 2017; Zhu, 2018). Generally, light energy capturing is much faster than electron transport in the thylakoid membranes; hence, over-excitation of the photosynthetic apparatus is a constant threat (Steyn, 2002).

Besides their antioxidant capacity, anthocyanins have the potential to reduce both the incidence and the severity of photo-oxidative damage. Due to the existence of these pigments in the vacuole apart from chloroplasts, therefore the light energy absorbed by them can't be transferred to chlorophyll to be used in photosynthesis (Gould, 2002). Which means that anthocyanins intercept a portion of supernumerary photons that would otherwise strike the chloroplasts, thus causing photo inhibition and increasing the ROS production and ROS-triggered damage (Gould, 2000). Accordingly, anthocyanins function as simple light filters, especially from the green – orange regions of the spectrum, and probably re-dissipate the absorbed energy as heat (Merzlyak and Chivkunova, 2000; Gould, 2002; Steyn, 2002; Landi, 2015; Trojak and Skowron, 2017).

Ultraviolet radiation-B (UV-B) stress: UV-B provokes ROS production in the living cells, which is very toxic because they react with vital biomolecules, altering their biological activities and causing oxidative damage. Anthocyanins, like other flavonoids, are well known for their ROS scavenging properties (Mahdavian, 2008; Goto, 2016). Moreover, since anthocyanins accumulate in the epidermal tissues of plants, they may contribute partially in preventing the penetration of UV-B to the photosynthetic mesophyll tissue (Mahdavian, 2008; Hatier and Gould, 2009; Landi, 2015). Numerous studies have indicated to anthocyanins increment under UV-B stress (Close and Beadle, 2003; Mahdavian, 2008; Tsurunaga, 2013; Goto, 2016; Li, 2020). UV-B induces a decline in anthocyanidin reductase (ANR) activity, which shifts the metabolic flux toward anthocyanin biosynthesis (Li, 2020). The common anthocyanin glycosides have negligible absorption in the UV region, but after acylation with phenolic acids, they gain the capacity to absorb parts of UV spectra (UV-A: 315–400 nm and UV-B: 280–315 nm) in addition to visible light (Hatier and Gould, 2009; Landi, 2015). However, UV filtering is not likely to be the main role of anthocyanins because acylated anthocyanins are not as common as the non-acylated forms in plant tissues (Hatier and Gould, 2009).

Anthocyanins and human health: Recently, anthocyanins have garnered a lot of attention for their potential preventative and /or therapeutic effects on health, including obesity prevention, cardiovascular diseases, antibacterial, anti-inflammatory and anticancer effects (Wu and Prior, 2005; Prior and Wu, 2006; Wallace and Giusti, 2015; Khoo, 2017).

Antibacterial effects: Previous studies referred to the antibacterial activity of anthocyanins against a wide range of microorganisms. This can be attributed to several mechanisms, such as destabilization and permeabilization of the plasma membrane (Burdulis, 2009), as well as induction cell damage and deformation by destroying the cell wall, membrane, and intercellular matrix integrity (Cisowska, 2011; Pojer, 2013; Khoo, 2017; Ma, 2019). Antibacterial property may also be related to the anti-adherence of bacteria to epithelial cells, which is essential for colonization and infection for many pathogens (Puupponen-Pimia, 2005).

Investigation of the antibacterial properties showed that anthocyanins-rich extracts inhibited the growth of a wide range of human pathogenic bacteria, both gram-negative and gram-positive (Puupponen-Pimia, 2005; Wu, 2008; Burdulis, 2009; Lacombe, 2010; Cote, 2011; Pagliarulo, 2015; Genskowsky, 2016; Aly, 2019). Although the inhibition is significantly more evident in gram-negative compared to gram-positive bacteria due to their structural variation (Puupponen-Pimia, 2001).

The cells of gram-negative bacteria are surrounded by an outer membrane, which acts as a preventive barrier against hydrophobic, but not hydrophilic compounds, due to the presence and features of lipopolysaccharide molecules in it (Puupponen-Pimia, 2005; Nohynek, 2006). These bacteria regulate the permeability of their cell through hydrophilic channels called porins, which allows nutrients to enter the cell cytoplasmic membrane (Nohynek, 2006).

Anti-inflammatory effects: Inflammation is the body's defence against stimuli. Usually, it is considered good for the body, but prolonged chronic inflammation can be harmful and result in different diseases such as obesity, type II diabetes, cardiovascular diseases and many types of cancer (Semaming, 2015; Yazhen, 2020).

Under acute inflammation, macrophages produce ROS while eliminating foreign particles and inducing inflammatory cytokines. ROS levels in the cell are balanced through detoxifying antioxidant enzymes (Medzhitov, 2010; Yazhen, 2020). Nevertheless, when the acute inflammation converts into chronic inflammation, the generation of ROS increases and becomes out of control, which will promote the inflammatory factors activated, aggravate the inflammatory response and gene mutations and finally lead to cancer. Many studies indicated that anthocyanins have strong anti-inflammatory activity in both in vivo and in vitro, and their effective mechanism may be the ability to scavenge ROS, reduce pro-inflammatory cytokines, and regulate antioxidants activity such as superoxide dismutase (Vendrame and Klimis, 2015; Pereira, 2017; Abdin, 2020; Yazhen, 2020).

Anticancer effects: The carcinogens can be induced by different factors, including pollutants and junk food. These carcinogens can be activated to produce free radicals that attack DNA and cause cancer. The anticancer property of anthocyanins is strongly related to their antioxidant capacity and cytotoxic action induction against cancer cells (Reddivari, 2007; Rugina, 2012; Diaconeasa, 2015; Thibado, 2018).

Pure anthocyanins and anthocyanin-rich extracts inhibit cell proliferation by blocking various stages of the cell cycle and their regulatory proteins (Wang and Stoner 2008). Interestingly, they can selectively inhibit the proliferation of cancer cells while having little influence on the proliferation of normal cells (Wang and Stoner 2008; Rugina, 2012; Bunea, 2013; Diaconeasa, 2017). Besides that, anthocyanins suppress angiogenesis, thereby inhibiting the growth and metastasis of tumours (Joshua, 2017; Tsakiroglou, 2019).

Besides, anthocyanins also can induce the apoptosis of cancer cells, which is not present in the tumour cells, through the internal mitochondrial pathway and the external death receptor pathway (Chang, 2005; Reddivari, 2007; Wang and Stoner 2008).

Anti-obesity effects: Obesity is characterized by an excessive accumulation of adipose tissue due to an imbalance between energy intake and expenditure. This condition is linked to lack of physical activity and unhealthy diet. Obesity may increase the risk of several diseases such as hypertension, heart disease and type II diabetes (Pojer, 2013; Jayarathne, 2019).

The role of anthocyanins in obesity is still controversial. However, previous studies indicated that anthocyanin consumption might aid in maintaining or reducing the bodyweight of obese healthy patients. In addition to improving lipid metabolism and energy balance (Bertoia, 2016; Azzini, 2017; Istek and Gurbuz, 2017; Sivamaruthi, 2020), inhibiting lipid absorption, suppressing food intake and regulating gut microbiota (Xie, 2018; Jayarathne, 2019).

Anti-diabetic effects: Diabetes mellitus is a metabolic syndrome caused by a combination of genetic and lifestyle patterns and arises when produced insulin is insufficient or inefficient. Diabetes leads to hyperglycemia, which in turn results in various short-term metabolic changes in lipid and protein metabolism and long-term irreversible vascular changes. Type II diabetes is commonly characterized by insulin resistance (reduced cell's sensitivity to insulin), an insulin deficiency, or both. Moreover, it occurs usually in adulthood to obese patients (Kosti and Kanakari, 2012).

Several studies demonstrated that anthocyanins can lower blood glucose levels via different mechanisms. In addition to their antioxidant capacity, anthocyanins can ameliorate insulin resistance, increase insulin secretion, improve liver function, and inhibit carbohydrate hydrolyzing enzymes (Prior and Wu, 2006; Belwal, 2017; Gowd, 2017).

Effects against cardiovascular diseases: As people age, the elastic fibres of arteries are oxidized and become harder, which represents a major cause of cardiovascular disease in the elderly (Yazhen, 2020).

The cardiovascular protective role of anthocyanins is strongly related to their properties against oxidative stress (Wallace, 2011). Besides, anthocyanins are well known for their capacity to decrease low-density lipoprotein cholesterol (LDL) (Pojer, 2013; Kruger, 2014; Lee, 2016; Reis, 2016), triglyceride (Reis, 2016) and blood pressure (Basu, 2010; Kruger, 2014; Reis, 2016; Igwe, 2017). Besides that, they inhibit platelet aggregation and activation (Saluk, 2012; Yang, 2012; Reis, 2016; Thompson, 2017). Anthocyanins normally reduce the risk of myocardial infarction (Cassidy, 2013) and inflammation in atherosclerosis (Reis, 2016). On the other hand, they improve high-density lipoprotein cholesterol (HDL) (Erlund, 2008; Hassellund, 2013; Kruger, 2014; Reis, 2016). Moreover, anthocyanin isolates and anthocyanin-rich mixtures of flavonoids may protect from lipid peroxidation, anti-inflammatory activity (Semaming, 2015), decreased capillary permeability and fragility, and membrane strengthening (Wallace, 2011).

Effects against neurodegenerative diseases: Neurodegenerative diseases are characterized by the loss of specific neuronal populations within the brain, brain stem, and spinal cord, resulting in significant cognitive and/or motor disorders (Radi, 2014).

The high oxygen consumption of the brain (20% more oxygen compared to mitochondrial respiratory tissues) increases the possibility of producing a large number of free radicals and makes the brain more vulnerable to oxidative stress. Taking into account that the neuron (the basic functional unit of the brain) contains a large number of polyunsaturated fatty acids, it can interact with ROS, leading to lipid peroxidation and apoptosis (Thakur, 2018).

Apoptosis is involved in many human diseases, including neurodegenerative disorders such as Alzheimer's disease (AD) and Parkinson's disease (PD). In these cases, the apoptotic rate increase causing tissue damage (Radi, 2014).

Diets rich in anthocyanins is associated with decreased risk of developing neurodegenerative diseases (Winter, 2017), through direct scavenging of ROS, increasing the activity of antioxidant enzymes (superoxide dismutase (SOD) and catalase (CAT)), elevating reduced glutathione GSH content, and reducing malondialdehyde MDA (Winter and Bickford, 2019; Yazhen, 2020).

Positive findings have been shown for the treatment of Alzheimer's disease with anthocyanin-rich extracts. These extracts can reduce the levels of the β -amyloid peptide ($A\beta$) and convert $A\beta$ aggregation to an alternate, non-toxic form (Yamakawa, 2016; Andrade Teles, 2018; Winter and Bickford, 2019), besides decrease tau phosphorylation induced by $A\beta$ (Badshah, 2015; Belkacemi and Ramassamy, 2016).

Moreover, anthocyanin consumption by old people at risk for dementia improves memory impairment via increasing neuronal signaling in brain centers mediating this function (Cho, 2003; Krikorian, 2010; Andrade Teles, 2018; Winter and Bickford, 2019; Yazhen, 2020).

Additionally, taking into consideration that anthocyanins are potent antioxidants, they can effectively prevent free radicals from damaging the dopamine-producing cells in the brain, thus inhibiting the development of Parkinson's disease (Kim, 2010; Strathearn, 2014; Qian, 2019). They may also alleviate neuro degeneration in Parkinson's disease by enhancing mitochondrial function (Strathearn, 2014).

Effects against ophthalmology: The blue light harm becomes greater with the increased usage of various electronic devices such as mobile phones, computers, and LED lights. The long term exposure to blue light irradiation to the retina can cause excessive free radical production. These free radicals can induce retinal pigment epithelial cell apoptosis, intraocular metabolic abnormalities and hindering the blood circulation, which in turn cause various dysfunctions such as myopia, cataracts, retinopathy, and other eye diseases (Yazhen, 2020).

The possible effective mechanism of anthocyanins is related to their scavenging ability for free radical, enhancing antioxidants activity, inhibiting intracellular calcium overload, in addition to the optic nerve protection (Yazhen, 2020), and suppression of retinal pigment epithelium apoptosis (Liu, 2012; Wang, 2015). Furthermore, anthocyanins have a relaxing effect on the ciliary muscle, which is important to treat myopia and glaucoma (Shim, 2012; Nomi, 2019).

It's worth noting that anthocyanins intake improves transient myopic shift, dark adaptation and the retinal blood circulation in normal-tension glaucoma patients (Yanamala, 2009; Ohguro, 2007; Nomi, 2019).

The Side Effects of Anthocyanins: Anthocyanins toxicity has not been shown in currently published human intervention studies (Wallace and Giusti, 2015). They are generally regarded as safe and well-tolerated in humans, taking into account the long history of consumption of foods-rich in such flavonoids (Corcoran, 2012).

Nevertheless, there is considerable evidence that such compounds are not risk-free (Lambert, 2007). A scientific expert team for the European Food Safety Authority considered the currently available toxicological studies for anthocyanins inadequate to determine a numerically acceptable daily intake (ADI) for them (European Food Safety Authority, 2013).

However, it is worth noting that the potential side effect of anthocyanin or any other flavonoid can be associated with high doses of intake even for a short duration (Azzini, 2017).

2. CONCLUSIONS

In the past decades, interest in dietary phenolic compounds, including anthocyanins, has increased substantially. Anthocyanins are phytopigments found in several plants. They are well known for their antioxidant activities, which is due to their unique structure. The studies summarized in this review demonstrate the importance of these compounds as natural colourants, plant stress markers and preventives for many human chronic diseases.

REFERENCES

Abdin M, Hamed YS, Akhtar HMS, Chen D, Chen G, Wan P, Zeng X, Antioxidant and anti-inflammatory activities of target anthocyanins di-glucosides isolated from *Syzygium cumini* pulp by high-speed counter-current chromatography, Journal of Food Biochemistry, 2020, e13209.

Ahmed NU, Park JI, Jung HJ, Hur Y, Nou IS, Anthocyanin biosynthesis for cold and freezing stress tolerance and desirable colour in *Brassica rapa*, Functional and Integrative Genomics, 15 (4), 2015, 383 - 394.

Aly AA, Ali HGM, Eliwa NER, Phytochemical screening, anthocyanins and antimicrobial activities in some berries fruits, *Journal of Food Measurement and Characterization*. 13, 2019, 911–920.

Andersen OM, Jordheim M, Anthocyanins, In: *Flavonoids: Chemistry, Biochemistry and Applications*, Andersen OM, Markham KR, CRC Press, Taylor & Francis Group, 2006, 471 - 552.

Andrade Teles RB, Diniz TC, Pinto TCC, Júnior RGO, Silva MG, Lavor EM, Fernandes AWC, Oliveira AP, Almeida Ribeiro FPR, Silva AAM, Cavalcante TCF, Junior LJQ, Silva Almeida JRG, Flavonoids as therapeutic agents in Alzheimer's and Parkinson's diseases: A systematic review of preclinical evidences, *Oxidative Medicine and Cellular Longevity*, 2018.

Azzini E, Giacometti J, Russo GL, Antiobesity effects of anthocyanins in preclinical and clinical studies, *Oxidative Medicine and Cellular Longevity*, 2017.

Azzini E, Venneria E, Ciarapica D, Foddai MS, Intorre F, Zaccaria M, Maiani F, Palomba L, Barnaba L, Tubili C, Maiani G, Polito A, Effect of red-orange juice consumption on body composition and nutritional status in overweight/obese female: A pilot study, *Oxidative Medicine and Cellular Longevity*, 2017.

Badshah H, Kim TH, Kim MO, Protective effects of anthocyanins against amyloid beta-induced neurotoxicity *in vivo* and *in vitro*, *Neurochemistry International*, 80, 2015, 51-59.

Basu A, Du M, Leyva MJ, Sanchez K, Betts NM, Wu M, Aston CE, Lyons TJ, Blueberries decrease cardiovascular risk factors in obese men and women with metabolic syndrome, *The Journal of Nutrition*, 140 (9), 2010, 1582 – 1587.

Belkacemi A, Ramassamy C, Innovative anthocyanin/anthocyanidin formulation protects SK-N-SH cells against the amyloid- β peptide-induced toxicity: relevance to Alzheimer's disease, *Central Nervous System Agents in Medicinal Chemistry*, 16, 2016, 37-49.

Belwal T, Nabavi SF, Nabavi SM, Habtemariam S, Dietary anthocyanins and insulin resistance: When food becomes a medicine, *Nutrients*, 9, 2017, 1111.

Bertoia ML, Rimm EB, Mukamal KJ, Hu FB, Willett WC, Cassidy A, Dietary flavonoid intake and weight maintenance: Three prospective cohorts of 124 086 US men and women followed for up to 24 years, *BMJ*, 352, 2016.

Bueno MJ, Saez-Plaza P, Ramos-Escudero F, Jiménez AM, Fett R, Asuero AG, Analysis and antioxidant capacity of anthocyanin pigments, Part II: chemical structure, color and intake of anthocyanins, *Critical Reviews in Analytical Chemistry*, 42, 2012, 126 – 151.

Bunea A, Rugina D, Scontta Z, Pop RM, Pinteá A, Socaciu C, Tabaran F, Grootaert C, Struijs K, VanCamp J, Anthocyanin determination in blueberry extracts from various cultivars and their anti-proliferative and apoptotic properties in B16-F10 metastatic murine melanoma cells, *Phytochemistry*, 95, 2013, 436-444.

Burdulis D, Sarkinas A, Jasutiene I, Stackevicene E, Nikolajevs L, Janulis V, Comparative study of anthocyanin composition, antimicrobial and antioxidant activity in bilberry (*Vaccinium myrtillus* L.) and blueberry (*Vaccinium corymbosum* L.) fruits, *Acta Poloniae Pharmaceutica - Drug Research*, 66 (4), 2009, 399–408.

Cassidy A, Mukamal KJ, Liu L, Franz M, Eliassen AH, Rimm EB, High anthocyanin intake is associated with a reduced risk of myocardial infarction in young and middle-aged women, *Circulation*, 127 (2), 2013, 188 – 196.

Chakovari SZ, Enteshari S, Qasimov N, Effect of salinity stress on biochemical parameters and growth of borage (*Borago officinalis* L.), *Iranian Journal of Plant Physiology*, 6 (2), 2015, 1673 – 1685.

Chalker-Scott L, Do anthocyanins function as osmo regulatory in leaf tissues?, *Advances in Botanical Research*, 37, 2002, 103-106.

Chang YC, Huang HP, Hsu JD, Yang SF, Wang CJ, *Hibiscus* anthocyanins rich extract induced apoptotic cell death in human promyelocytic leukaemia cells, *Toxicology and Applied Pharmacology*, 205, 2005, 201-212.

Chanoca A, Kovinich N, Burkel B, Stecha S, Bohorquez-Restrepo A, Ueda T, Eliceiri KW, Grotewold E, Otegui MS, Anthocyanin vacuolar inclusions form by a micro autophagy mechanism, *The Plant Cell*, 27, 2015, 2545 – 2559.

Chemler JA, Leonard E, Koffas MAG, Flavonoid Bio transformations in Microorganisms, In *Anthocyanins: Biosynthesis, Functions, and Applications*, Gould K, Davies K, Winefield C, Springer, 2009, 191 – 238.

Cho J, Kang JS, Long PH, Jing J, Back Y, Chung KS, Antioxidant and memory-enhancing effects of purple sweet potato anthocyanin and cordyceps mushroom extract, *Archives of Pharmacal Research*, 26 (10), 2003, 821-825.

Chu WK, Cheung SCM, Lau RAW, Benzie IFF, Bilberry (*Vaccinium myrtillus* L.), In *Herbal Medicine: Bio molecular and Clinical Aspects*, 2nd edition, Benzie, I. F. F. and Wachtel- Galor, S, CRC Press, 2011, 55 -72.

Chunthaburee S, Sakuanrungrsirikul S, Wongwarat T, Sanitchon J, Pattanagul W, Theerakulpisut P, Changes in anthocyanin content and expression of anthocyanin synthesis genes in seedlings of black glutinous rice in response to salt stress, *Asian Journal of Plant Science*, 15 (3-4), 2016, 56 - 65.

Cisowska A, Wojnicz D, Hendrich AB, Anthocyanins as antimicrobial agents of natural plant origin, *Natural Product Communications*, 6 (1), 2011, 149-156.

Close DC, Beadle CL, The Eco physiology of foliar anthocyanin, *The Botanical Review*, 69 (2), 2003, 149 – 161.

Code of Federal Regulations, 21 CFR Part 73: Color Additives Exempt from Certification, Washington, DC, U.S. Food Drug Admin, 2020.

Corcoran MP, McKay DL, Blumberg JB, Flavonoid basics: Chemistry, sources, mechanisms of action, and safety, *Journal of nutrition in gerontology and geriatrics*, 31 (3), 2012, 176-189.

Cote J, Caillet S, Doyon G, Dussault D, Sylvain JF, Lacroix M, Antimicrobial effect of cranberry juice and extracts, *Food Control*, 22, 2011, 1413-1418.

Dai LP, Dong XJ, Ma HH, Antioxidative and chelating properties of anthocyanins in *Azolla imbricate* induced by cadmium, *Polish Journal of Environmental Studies*, 21 (4), 2012, 837- 844.

Delgado-Vargas F, Jimenez AR, Paredes-Lopez O, Natural Pigments: Carotenoids, Anthocyanins, and Betalains — Characteristics, Biosynthesis, Processing, and Stability, *Critical Reviews in Food Science and Nutrition*, 40 (3), 2000, 173 – 289.

Deroles S, Anthocyanin Biosynthesis in Plant Cell Cultures: A Potential Source of Natural Colourants, In *Anthocyanins: Biosynthesis, Functions, and Applications*, Gould K, Davies K, Winefield C, Springer, 2009, 107 – 167.

Diaconeasa Z, Ayvaz H, Rugina D, Leopold L, Stanila A, Socaciu C, Tabaran F, Luput L, Mada DC, Pintea A, Jefferson A, Melanoma inhibition by anthocyanins is associated with the reduction of oxidative stress biomarkers and changes in mitochondrial membrane potential, *Plant Foods for Human Nutrition*, 72, 2017, 404–410.

Diaconeasa Z, Leopold L, Rugina D, Ayvaz H, Socaciu C, Anti proliferative and antioxidant properties of anthocyanin-rich extracts from blueberry and blackcurrant juice, *International Journal of Molecular Sciences*, 16, 2015, 2352-2365.

Erlund I, Koli R, Alfthan G, Marniemi J, Puukka P, Mustonen P, Mattila P, Jula A, Favorable effects of berry consumption on platelet function, blood pressure, and HDL cholesterol, *American Journal of Clinical Nutrition*, 87, 2008, 323–331.

Eryilmaz F, The relationships between salt stress and anthocyanin content in higher plants, *Biotechnology & Biotechnological Equipment*, 20 (1), 2006, 47 - 52.

European Food Safety Authority, Scientific Opinion on the re-evaluation of anthocyanins (E 163) as a food additive, *EFSA Journal*, 11 (4), 2013, 3145.

Fang J, Classification of fruits based on anthocyanin types and relevance to their health effects, *Nutrition*, 31, 2015, 1301 – 1306.

Gaulejac NS, Glories Y, Vivas N, Free radical scavenging effect of anthocyanins in red wines, *Food Research International*, 32, 1999, 327 - 333.

Genskowsky E, Puente LA, Pérez-Alvarez JA, Lopez JF, Munoz LA, Viuda-Martos M, Determination of poly phenolic profile, antioxidant activity and antibacterial properties of maqui [*Aristotelia chilensis* (Molina) Stuntz] a Chilean blackberry, *J. Sci. Food Agric*, 96, 2016, 4235–4242.

Giusti MM, Wrolstad RE, Acylated anthocyanins from edible sources and their applications in food systems, *Biochemical Engineering Journal*, 14, 2003, 217–225.

Goto E, Hayashi K, Furuyama S, Hikosaka S, Ishigami Y, Effect of UV light on phytochemical accumulation and expression of anthocyanin biosynthesis genes in red leaf lettuce, *ISHS Acta Horticulturae*, 1134, 2016.

Gould KS, Markham KR, Smith RH, Goris JJ, Functional role of anthocyanins in the leaves of *Quintinia serrate* A. Cunn, Journal of Experimental Botany, 51 (347), 2000, 1107 – 1115.

Gould KS, Neill SO, Vogelmann TC, A unified explanation for anthocyanins in leaves?, Advances in Botanical Research, 37, 2002, 167 – 192.

Gowd V, Jia Z, Chen W, Anthocyanins as promising molecules and dietary bioactive components against diabetes-A review of recent advances, Trends in Food Science & Technology, 68, 2017, 1-13.

Grotewold E, Davies K, Trafficking and sequestration of anthocyanins, Natural Product Communications, 3 (8), 2008, 1251 – 1258.

Harborne JB, Williams CA, Anthocyanins and other flavonoids, Natural Product Reports, 18, 2001, 310 – 333.

Hassellund SS, Flaa A, Kjeldsen SE, Seljeflot I, Karlsen A, Erlund I, Rostrup M, Effects of anthocyanins on cardiovascular risk factors and inflammation in pre-hypertensive men: a double-blind randomized placebo-controlled crossover study, Journal of Human Hypertension, 27, 2013, 100 – 106.

Hatier JHB, Gould KS, Anthocyanin function in vegetative organs, In Anthocyanins: Biosynthesis, Functions, and Applications, Gould K, Davies K, Winefield C, Springer, 2009, 1 – 19.

He Q, Ren Y, Zhao W, Li R, Zhang L, Low temperature promotes anthocyanin biosynthesis and related gene expression in the seedlings of purple head Chinese cabbage (*Brassica rapa* L.), Genes, 11, 2020, 81.

Hirsch GE, Martins LAM, Anthocyanins: chemical features, food sources and health benefits, In Handbook of anthocyanins food sources, chemical applications and health benefits, Warner LM, Nova Science Publishers, 2015, 227 – 248.

Igwe EO, Charlton KE, Roodenrys S, Kent K, Fanning K, Netzel ME, Anthocyanin-rich plum juice reduces ambulatory blood pressure but not acute cognitive function in younger and older adults: A pilot cross-over dose-timing study, Nutrition Research, 47, 2017, 28 – 43.

Iseri OD, Korpe DA, Sahin FI, Haberal M, High salt-induced oxidative damage and antioxidant response in tomato grafted on tobacco, Chilean Journal of Agricultural Research, 75 (2), 2015, 192 – 201.

Istek N, Gurbuz O, Investigation of the impact of blueberries on metabolic factors influencing health, Journal of Functional Foods, 38, 2017, 298–307.

Jayarathne S, Stull AJ, Park OH, Kim JH, Thompson L, Moustaid-Moussa N, Protective effects of anthocyanins in obesity-associated inflammation and changes in the gut microbiome, Molecular Nutrition & Food Research, 63, 2019.

Joshua M, Okere C, Sylvester O, Yahaya M, Precious O, Dluya T, Um J, Neksumi M, Boyd J, Vincent-Tyndall J, Choo D, Gutsaeva DR, Jahng WJ, Disruption of angiogenesis by anthocyanin-rich extracts of *Hibiscus sabdariffa*, International Journal of Scientific and Engineering Research, 8 (2), 2017, 299–307.

Kebbas S, Benseddik T, Makhloufi H, Aid F, Physiological and biochemical behaviour of *Gleditsia triacanthos* L. young seedlings under drought stress conditions, Not Bot Horti Agrobo, 46 (2) 2018, 585-592.

Khoo HE, Azlan A, Tang ST, Lim SM, Anthocyanidins and anthocyanins: coloured pigments as food, pharmaceutical ingredients, and the potential health benefits, Food & Nutrition Research, 61 (1), 2017, 1361779.

Kielkowska A, Grzebelus E, Lis-Krzyscin A, Mackowska K, Application of the salt stress to the protoplast cultures of the carrot (*Daucus carota* L.) and evaluation of the response of regenerants to soil salinity, Plant Cell, Tissue and Organ Culture (PCTOC), 137, 2019, 379 – 395.

Kim HG, Ju MS, Shim JS, Kim MC, Lee SH, Huh Y, Kim SY, Oh MS, Mulberry fruit protects dopaminergic neurons in toxin-induced Parkinson's disease models, British Journal of Nutrition, 104, 2010, 8–16.

Kosti M, Kanakari M, Education and diabetes mellitus, Health Science Journal, 6 (4), 2012, 654-662.

Kovinich N, Kayanja G, Chanoca A, Otegui MS, Grotewold E, Abiotic stresses induce different localizations of anthocyanins in *Arabidopsis*, Plant Signaling & Behavior, 10, 2015,7.

Krikorian R, Shidler MD, Nash TA, Kalt W, Vinqvist-Tymchuk MR, Shukitt-Hale BS, Joseph JA, Blueberry supplementation improves memory in older adults, Journal of Agricultural and Food Chemistry, 58 (7), 2010, 3996–4000.

Kruger MJ, Davies N, Myburgh KH, Lecour S, Pro anthocyanidins, anthocyanins and cardiovascular diseases, *Food Research International*, 59, 2014, 41 – 52.

Lacombe A, Wu, VCH, Tyler, S, Edwards K, Antimicrobial action of the American cranberry constituents; phenolics, anthocyanins, and organic acids, against *Escherichia coli* O157:H7, *International Journal of Food Microbiology*, 139 (1–2), 2010, 102–107.

Lambert JD, Sang S, Yang CS, Possible controversy over dietary polyphenols: benefits vs. risks, *Chemical Research in Toxicology*, 20, 2007, 583–585.

Landi M, Tattini M, Gould KS, Multiple functional roles of anthocyanins in plant-environment interactions, *Environmental and Experimental Botany*, 119, 2015, 4 – 17.

Latti AK, Kainulainen PS, Hayirlioglu-Ayaz S, Ayaz FA, Riihinen KR, Characterization of anthocyanins in caucasian blueberries (*Vaccinium arctostaphylos* L.) native to Turkey, *Journal of Agricultural and Food Chemistry*, 57, 2009, 5244 – 5249.

Lee M, Sorn SR, Park Y, Park H, Anthocyanin rich-black soybean test improved visceral fat and plasma lipid profiles in overweight/obese Korean adults: A randomized controlled trial, *Journal of Medicinal Food*, 19 (11), 2016, 1–9.

Li W, Tan L, Zou Y, Tan X, Huang J, Chen W, Tang Q, The Effects of ultraviolet A/B treatments on anthocyanin accumulation and gene expression in dark-purple tea cultivar ‘Ziyan’ (*Camellia sinensis*), *Molecules*, 25, 2020, 354.

Liu Y, Song X, Zhang X, Zhou F, Wang D, Wei Y, Gao F, Xie L, Jia G, Wu W, Ji B, Blueberry anthocyanins: protection against ageing and light-induced damage in retinal pigment epithelial cells, *British Journal of Nutrition*, 108, 2012, 16–27.

Liu Y, Tikunov Y, Schouten RE, Marcelis LFM, Visser RGF, Bovy A, Anthocyanin biosynthesis and degradation mechanisms in Solanaceous vegetables: A review, *Frontiers in Chemistry*, 6, 2018, 52.

Ma Y, Ding S, Fei Y, Liu G, Jang H, Fang J, Antimicrobial activity of anthocyanins and catechins against foodborne pathogens *Escherichia coli* and *Salmonella*, *Food Control*, 106, 2019, 106712.

Mahdavian K, Ghorbanli M, Kalantari KM, The effects of ultraviolet radiation on the contents of chlorophyll, flavonoid, anthocyanin and proline in *Capsicum annum* L, *Turkish Journal of Botany*, 32, 2008, 25 – 33.

Manetas Y, Why some leaves are anthocyanins and why most anthocyanins leaves are red?, *Flora*, 201 (3), 2006, 163–177.

Martin J, Navas MJ, Jimenez-Moreno AM, Asuero AG, Anthocyanin Pigments: Importance, Sample Preparation and Extraction, In *Phenolic Compounds - Natural Sources, Importance and Applications*, Soto- Hernandez M, Palma-Tenango M, Garcia-Mateos R, Books on Demand, 2017, 117 – 152.

Medzhitov R, Inflammation: new adventures of an old flame, *Cell*, 140 (6), 2010, 771–776.

Merzlyak MN, Chivkunova OB, Light-stress-induced pigment changes and evidence for anthocyanin photo protection in apples, *Journal of Photochemistry and Photobiology B. Biology*, 55, 2000, 155–163.

Mortensen A, Carotenoids and other pigments as natural colourants, *Pure and Applied Chemistry*, 78 (8), 2006, 1477–1491.

Nakabayashi R, Mori T, Saito K, Alternation of flavonoid accumulation under drought stress in *Arabidopsis thaliana*, *Plant Signaling and Behavior*, 9 (8), 2014, e29518.

Nakabayashi R, Yonekura-Sakakibara K, Urano K, Suzuki M, Yamada Y, Nishizawa T, Matsuda F, Kojima M, Sakakibara H, Shinozaki K, Michael AJ, Tohge T, Yamazaki M, Saito K, Enhancement of oxidative and drought tolerance in *Arabidopsis* by over accumulation of antioxidant flavonoids, *The Plant Journal*, 77, 2014, 367–379.

Nohynek LJ, Alakomi HL, Kahkonen MP, Heinonen M, Helander KM, Oksman- Caldentey KM, Puupponen-Pimia RH, Berry phenolics: antimicrobial properties and mechanisms of action against severe human pathogens, *Nutrition and Cancer*, 54, 2006, 18–32.

Nomi Y, Kurashige KI, Matsumoto H, Therapeutic effects of anthocyanins for vision and eye health, *Molecules*, 24, 2019, 3311.

Ohguro I, Ohguro H, Nakazawa M, Effects of anthocyanins in black currant on retinal blood flow circulation of patients with normal-tension glaucoma. A pilot study, *Hirosaki Medical Journal*, 59 (1), 2007, 23–32.

Oku M, Sakai Y, Three distinct types of micro autophagy based on membrane dynamics and molecular machinery, *Bio Essays*, 40, 2018.

Omidi FJ, Shoja HM, Sariri R, Effect of water-deficit stress on secondary metabolites of *Melissa officinalis* L.: role of exogenous salicylic acid, *Caspian J. Environ. Sci.*, 16 (2), 2018, 121-134.

Pagliarulo C, De Vito V, Picariello G, Colicchio R, Pastore G, Salvatore P, Volpe MG, Inhibitory effect of pomegranate (*Punica granatum* L.) polyphenol extracts on the bacterial growth and survival of clinical isolates of pathogenic *Staphylococcus aureus* and *Escherichia coli*, *Food Chemistry*, 2015.

Passeri V, Koes R, Quattrocchio FM, New challenges for the design of high-value plant products: Stabilization of anthocyanins in plant vacuoles, *Frontiers in Plant Science*, 7, 2016, 153.

Pereira DM, Valentao P, Pereira JA, Andrade PB, Phenolics: from chemistry to biology, *Molecules*, 14, 2009, 2202 - 2211.

Pereira SR, Pereira R, Figueiredo I, Freitas V, Dinis TCP, Almeida LM, Comparison of anti-inflammatory activities of an anthocyanin-rich fraction from Portuguese blueberries (*Vaccinium corymbosum* L.) and 5-aminosalicylic acid in a TNBS-induced colitis rat model, *PLOS ONE*, 12 (3), 2017, e0174116.

Pojer E, Mattivi F, Johnson D, Stockley CS, The case for anthocyanin consumption to promote human health: A review, *Comprehensive Reviews in Food Science and Food Safety*, 12, 2013, 483 - 508.

Prior RL, Wu X, Anthocyanins: Structural characteristics that result in unique metabolic patterns and biological activities, *Free Radical Research*, 40 (10), 2006, 1014 – 1028.

Puupponen-Pimia R, Nohynek L, Alakomi HA, Oksman-Caldentey KM, The action of berry phenolics against human intestinal pathogens, *Bio Factors*, 23, 2005, 243–251.

Puupponen-Pimia R, Nohynek L, Hartmann-Schmidlin S, Kahkonen M, Heinonen M, Maatta-Riihinen K, Oksman-Caldentey KM, Berry phenolics selectively inhibit the growth of intestinal pathogens, *Journal of Applied Microbiology*, 98, 2005, 991–1000.

Puupponen-Pimia R, Nohynek L, Meier C, Kahkonen M, Heinonen M, Hopia A, Oksman-Caldentey KM, Antimicrobial properties of phenolic compounds from berries, *Journal of Applied Microbiology*, 90 (4), 2001, 494–507.

Qian F, Wang M, Wang J, Lu C, Anthocyanin-rich blueberry extract ameliorates the behavioral deficits of MPTP-induced mouse model of Parkinson's disease via anti-oxidative mechanisms, *Yangtze Medicine*, 3, 2019, 72-78.

Radi E, Formichi P, Battisti C, Federico A, Apoptosis and oxidative stress in neurodegenerative diseases, *Journal of Alzheimer's Disease*, 42 (3), 2014, S125–S152.

Reddivari L, Vanamala J, Chintharlapalli S, Safe SH, Miller JC, Anthocyanin fraction from potato extracts is cytotoxic to prostate cancer cells through activation of caspase-dependent and caspase-independent pathways, *Carcinogenesis*, 28, 2007, 2227-2235.

Reis JF, Monteiro VVS, Gomes RS, do Carmo MM, da Costa GV, Ribera PC, Monteiro MC, Action mechanism and cardiovascular effect of anthocyanins: a systematic review of animal and human studies, *Journal of Translational Medicine*, 14, 2016, 315.

Rugina D, Sconta Z, Leopold L, Pintea A, Bunea A, Socaciu C, Antioxidant activities of chokeberry extracts and the cytotoxic action of their anthocyanin fraction on HeLa human cervical tumour cells, *Journal of Medicinal Food*, 15, 2012, 700-706.

Sadilova E, Stintzing FC, Carle R, Anthocyanins, colour and antioxidant properties of eggplant (*Solanum melongena* L.) and violet pepper (*Capsicum annuum* L.) peel extracts, *Z. Naturforsch. C*, 61, 2006, 527–535.

Saluk J, Bijak M, Kolodziejczyk-Czepas J, Posmyk MM, Janas KM, Wachowicz B, Anthocyanins from red cabbage extract – evidence of protective effects on blood platelets, *Central European Journal of Biology*, 7 (4), 2012, 655 – 663.

Santos-Buelga C, Mateus N, De Freitas V, Anthocyanins. Plant pigments and beyond, *Journal of Agricultural and Food Chemistry*, 2014.

Schulz E, Tohge T, Zuther E, Fernie AR, Hinch DK, Flavonoids are determinants of freezing tolerance and cold acclimation in *Arabidopsis thaliana*, *Scientific Reports*, 6, 2016, 34027.

Schulz E, Tohge T, Zuther E, Fernie AR, Hinch DK, Natural variation in flavonol and anthocyanin metabolism during cold acclimation in *Arabidopsis thaliana* accessions, *Plant, Cell and Environment*, 38, 2015, 1658–1672.

Semaming Y, Pannengpetch P, Chattipakorn SC, Chattipakorn N, Pharmacological properties of proto catechuic acid and its potential roles as complementary medicine, *Evidence-Based Complementary and Alternative Medicine*, 2015.

Shim SH, Kim JM, Choi CY, Kim CY, Park KH, *Ginkgo biloba* extract and bilberry anthocyanins improve visual function in patients with normal-tension glaucoma, *Journal of Medicinal Food*, 15 (9), 2012, 818–823.

Shoeva OY, Gordeeva EI, Arbuzova VS, Khlestkina EK, Anthocyanins participate in the protection of wheat seedlings from osmotic stress, *Cereal Research Communications*, 45 (1), 2017, 47–56.

Sivamaruthi BS, Kesika P, Chaiyasut C, The influence of supplementation of anthocyanins on obesity-associated comorbidities: A concise review, *Foods*, 9, 2020, 687.

Sperdoulis I, Moustakas M, Interaction of proline, sugars, and anthocyanins during photosynthetic acclimation of *Arabidopsis thaliana* to drought stress, *Journal of Plant Physiology*, 169, 2012, 577–585.

Steyn WJ, Wand SJE, Holcroft DM, Jacobs G, Anthocyanins in vegetative tissues: A proposed unified function in photo protection, *New Phytologist*, 155, 2002, 349 – 361.

Strathearn KE, Yousef GG, Grace MH, Roy SL, Tambe MA, Ferruzzi MG, Wu QL, Simon JE, Lila MA, Rochet JC, Neuro protective effects of anthocyanin and pro anthocyanidin-rich extracts in cellular models of Parkinson's disease, *Brain Research*, 1555, 2014, 60-77.

Tanaka Y, Sasaki N, Ohmiya A, Biosynthesis of plant pigments: Anthocyanins, betalains and carotenoids, *The Plant Journal*, 54, 2008, 733 – 749.

Thakur AK, Kamboj P, Goswami K, Ahuja K, Pathophysiology and management of Alzheimer's disease: An overview, *Journal of Analytical & Pharmaceutical Research*, 9 (2), 2018, 226–235.

Thibado SP, Thornthwaite JT, Ballard TK, Goodman BT, Anticancer effects of Bilberry anthocyanins compared with Nutra Nano Sphere encapsulated bilberry anthocyanins, *Molecular and Clinical Oncology*, 8, 2018, 330-335.

Thompson K, Hosking H, Pederick W, Singh I, Santhakumar AB, The effect of anthocyanin supplementation in modulating platelet function in sedentary population: a randomized, double-blind, placebo-controlled, cross-over trial, *British Journal of Nutrition*, 118, 2017, 368 – 374.

Tomic J, Stajic ZK, Pesakovic M, The role of berry growing technology on bioactive compound improvement, In *Phenolic compounds: structure, uses and health benefits*, Oliver P, Villem A, Nova Science Publishers, 2017, 37-92.

Trojak M, Skowron E, Role of anthocyanins in high-light stress response, *World Science News*, 81 (2), 2017, 150-168.

Tsakiroglou P, VandenAkker NE, Del Bo C, Riso P, Klimis-Zacas D, Role of berry anthocyanins and phenolic acids on cell migration and angiogenesis: An updated overview, *Nutrients*, 11, 2019, 1075.

Tsurunaga Y, Takahashi T, Katsube T, Kudo A, Kuramitsu O, Ishiwata M, Matsumoto S, Effects of UV-B irradiation on the levels of anthocyanin, rutin and radical scavenging activity of buckwheat sprouts, *Food Chemistry*, 141, 2013, 552–556.

Ubi BE, Honda C, Bessho H, Kondo S, Wada M, Kobayashi S, Moriguchi T, Expression analysis of anthocyanin biosynthetic genes in apple skin: effect of UV-B and temperature, *Plant Science*, 170, 2006, 571–578.

Usenik V, Stampar F, Veberic R, Anthocyanins and fruit colour in plums (*Prunus domestica* L.) during ripening, *Food Chemistry*, 114, 2009, 529–534.

Vendrame S, Klimis D, Anti-inflammatory effect of anthocyanins via modulation of nuclear factor- B and mitogen-activated protein kinase signaling cascades, *Nutrition Reviews*, 73 (6), 2015, 348-58.

Vermerris W, Nicholson R, *Phenolic compound biochemistry*, Netherlands, Springer, 2006, 1-62.

Wallace TC, Anthocyanins in cardiovascular disease, *Advances in Nutrition*, 2 (1), 2011, 1–7.

Wallace TC, Giusti MM, Anthocyanins—Nature's bold, beautiful, and health-promoting colours, *Foods*, 8, 2019, 550.

Wallace TC, Giusti MM, Anthocyanins, *Advances in Nutrition*, 6 (5), 2015, 620–622.

- Wang LS, Stoner GD, Anthocyanins and their role in cancer prevention, *Cancer Lett.*, 269 (2), 2008, 281–90.
- Wang Y, Zhao L, Lu F, Yang X, Deng Q, Ji B, Huang F, Retino protective effects of bilberry anthocyanins via antioxidant, anti-inflammatory, and anti-apoptotic mechanisms in a visible-light-induced retinal degeneration model in pigmented rabbits, *Molecules*, 20, 2015, 22395–22410.
- Winter AN, Bickford PC, Anthocyanins and their metabolites as therapeutic agents for neurodegenerative disease, *Antioxidants*, 8, 2019, 333.
- Winter AN, Brenner MC, Punessen N, Snodgrass M, Byars C, Arora Y, Linseman DA, Comparison of the neuro protective and anti-inflammatory effects of the anthocyanin metabolites, proto catechuic acid and 4-hydroxybenzoic acid, *Oxidative Medicine and Cellular Longevity*, 2017.
- Wrolstad RE, Anthocyanin pigments-bioactivity and colouring properties, *Journal of Food Science*, 69 (5), 2004, C419–C425.
- Wrolstad RE, Culver CA, Alternatives to those artificial FD&C food colourants, *Annual Review of Food Science and Technology*, 3, 2012, 59–77.
- Wu VC, Qiu X, Bushway A, Harper L, Antibacterial effects of American cranberry (*Vaccinium macrocarpon*) concentrate on foodborne pathogens, *LWT - Food Science and Technology*, 41, 2008, 1834-1841.
- Wu X, Prior RL, Systematic identification and characterization of anthocyanins by HPLC-ESI-MS/MS in common foods in the United States: Fruits and berries, *Journal of Agricultural and Food Chemistry*, 53, 2005, 2589 - 2599.
- Xie L, Su H, Sun C, Zheng X, Chen W, Recent advances in understanding the anti-obesity activity of anthocyanins and their biosynthesis in microorganisms, *Trends in Food Science & Technology*, 72, 2018, 13–24.
- Yamakawa MY, Uchino K, Watanabe Y, Adachi T, Nakanishi M, Ichino H, Hongo K, Mizobata T, Kobayashi S, Nakashima K, Kawata Y, Anthocyanin suppresses the toxicity of A β deposits through the diversion of molecular forms *in vitro* and *in vivo* models of Alzheimer's disease, *Nutritional Neuroscience*, 19 (1), 2016, 32-42.
- Yanamala N, Tirupula KC, Balem F, Klein-Seetharaman J, pH-dependent interaction of rhodopsin with cyanidin-3-glucoside, *Structural aspects, Photochemistry and Photobiology*, 85, 2009, 454–462.
- Yang Y, Shi Z, Rehemana A, Jin JW, Li C, Wang Y, Andrews MC, Chen P, Zhu G, Ling W, Ni H, Plant food delphinidin-3-glucoside significantly inhibits platelet activation and thrombosis: novel protective roles against cardiovascular diseases, *PLoS One*, 7 (5), 2012, e37323.
- Yazhen S, Wenju W, Panpan Z, Yuanyuan Y, Panpan D, Wusen Z, Yanling W, Anthocyanins: Novel Antioxidants in diseases prevention and human health, In Badria, F.A. and Ananga, A. *Flavonoids - A Coloring Model for Cheering up Life*, 2020.
- Zhang KM, Yua HJ, Shia K, Zhou YH, Yu JQ, Xia XJ, Photo protective roles of anthocyanins in *Begonia semperflorens*, *Plant Science*, 179, 2010, 202–208.
- Zhang O, Zhai J, Shao L, Lin W, Peng C, Accumulation of anthocyanins: An adaptation strategy of *Mikania micrantha* to low temperature in winter, *Frontiers in Plant Science*, 10, 2019.
- Zhao Z, Yan H, Zheng R, Saeed KM, Fu X, Tao Z, Zhang Z, Anthocyanins characterization and antioxidant activities of sugarcane (*Saccharum officinarum* L.) rind extracts, *Industrial Crops and Products*, 113, 2018, 38–45.
- Zhu H, Zhang TJ, Zheng J, Huang XD, Yu ZC, Peng CL, Chow WS, Anthocyanins function as a light attenuator to compensate for insufficient photo protection mediated by non-photochemical quenching in young leaves of *Acmena acuminatissima* in winter, *Photosynthetica*, 56 (1), 2018, 445-454.